CONTENTS

Symposium on Biomonitoring and Susceptibility Markers in Human Cancer: Applications in Molecular Epidemiology and Risk Assessment. Part II. Exposure Monitoring and Molecular Dosimetry

Preface 3

Assessment of Carcinogen Exposure and the Use of Other Risk Markers in Populations

F. A. Beland and M. C. Poirier. Significance of DNA adduct studies in animal models for cancer molecular dosimetry and risk asessment 5

V. E. Walker, T. R. Fennell, P. B. Upton, J. P. MacNeela, and J. A. Swenberg. Molecular dosimetry of DNA and hemoglobin adducts in mice and rats exposed to ethylene oxide 11

P. B. Farmer, E. Bailey, S. Naylor, D. Anderson, A. Brooks, J. Cushnir, J. H. Lamb, O. Sepai, and Y-S. Tang. Identification of endogenous electrophiles by means of mass spectrometric determination of protein and DNA adducts 19

J. H. M. van Delft, M. J. M. van Winden, A. M. C. van den Ende, and R. A. Baan. Determining N7-alkyguanine adducts by immunochemical methods and HPLC with electrochemical detection: applications in animal studies and in monitoring human exposure to alkylating agents 25

D. E. G. Shuker, V. Prevost, M. D. Friesen, D. Lin, H. Ohshima, and H. Bartsch. Urinary markers for measuring exposure to endogenous and exogenous alkylating agents and precursors 33

M. Törnqvist and A. Kautiainen. Adducted proteins for identification of endogenous electrophiles 39

D. H. Phillips and A. Hewer. DNA adducts in human urinary bladder and other tissues 45

S. R. Tannenbaum, P. L. Skipper, J. S. Wishnok, W. G. Stillwell, B. W. Day, and K. Taghizadeh. Characterization of various classes of protein adducts 51

S. S. Hecht, S. G. Carmella, P. G. Foiles, S. E. Murphy, and L. A. Peterson. Tobacco-specific nitrosamine adducts: studies in laboratory animals and humans 57

H.-G. Neumann, G. Birner, P. Kowallik, D. Schütze, and I. Zwirner-Baier. Hemoglobin adducts of *N*-substituted aryl compounds in exposure control and risk assessment 65

E. Kriek, F. J. Van Schooten, M. J. X. Hillebrand, F. E. Van Leeuwen, L. Den Engelse, A. J. A. De Looff, and A. P. G. Dijkmans. DNA adducts as a measure of lung cancer risk in humans exposed to polycyclic aromatic hydrocarbons 71

E. Grzybowska, K. Hemminki, and M. Chorąży. Seasonal variations in levels of DNA adducts and X-spots in human populations living in different parts of Poland 77

J. L. Mumford, X. Lee, J. Lewtas, T. L. Young, and R. M. Santella. DNA adducts as biomarkers for assessing exposure to polycyclic aromatic hydrocarbons in tissues from Xuan Wei women with high exposure to coal combustion emissions and high lung cancer mortality 83

J. Lewtas, J. Mumford, R. B. Everson, B. Hulka, T. Wilcosky, W. Kozumbo, C. Thompson, M. George, L. Dobiáš, R. Šrám, X. Li, and J. Gallagher. Comparison of DNA adducts from exposure to complex mixtures in various human tissues and experimental systems 89

J. C. Harrison, Magda Carvajal, and R. C. Garner. Does aflatoxin exposure in the United Kingdom constitute a cancer risk? 99

J. D. Groopman, C. P. Wild, J. Hasler, C. Junshi, G. N. Wogan, and T. W. Kensler. Molecular epidemiology of aflatoxin exposures: validation of aflatoxin-N7-guanine levels in urine as a biomarker in experimental rat models and humans 107

C. P. Wild, L. A. M. Jansen, L. Cova, and R. Montesano. Molecular dosimetry of aflatoxin exposure: contribution to understanding the multifactorial etiopathogenesis of primary hepatocellular carcinoma with particular reference to hepatitis B virus 115

R. J. Turesky, W. G. S. Stillwell, P. L. Skipper, and S. R. Tannenbaum. Metabolism of the food-borne carcinogens 2-amino-3-methylimidazo [4,5-*f*]quinoline and 2-amino-3,8-dimethylimidazo [4,5-*f*]quinoxaline in the rat as a model for human biomonitoring 123

K. Wakabayashi, H. Ushiyama, M. Takahashi, H. Nukaya, S-B. Kim, M. Hirose, M. Ochiai, T. Sugimura, and M. Nagao. Exposure to heterocyclic amines 129

R. J. Albertini, J. A. Nicklas, J. C. Fuscoe, T. R. Skopek, R. F. Branda, and J. P. O'Neill. *In vivo* mutations in human blood cells: biomarkers for molecular epidemiology 135

S. A. Kyrtopoulos, V. L. Souliotis, C. Valavanis, V. A. Boussiotis, and G. A. Pangalis. Accumulation of O^6-methylguanine in human DNA after therapeutic exposure to methylating agents and its relationship with biological effects 143

M. C. Poirier, E. Reed, H. Shamkhani, R. E. Tarone, and S. Gupta-Burt. Platinum drug–DNA interactions in human tissues measured by cisplatin–DNA enzyme–linked immunosorbent assay and atomic absorbance spectroscopy 149

J. S. Wishnok, S. R. Tannenbaum, W. G. Stillwell, J. A. Glogowski, and C. D. Leaf. Urinary markers for exposures to alkylating or nitrosating agents 155

I. O'Neill, S. Bingham, A. Ellul, and B. Incaurgarat. Magnetic microcapsule exploration in the gastrointestinal cavity of the origins of colorectal cancer-associated DNA-damaging agents in the human diet 161

A. Scalmati and M. Lipkin. Proliferation and differentiation biomarkers in colorectal mucosa and their application to chemoprevention studies 169

Manuscripts from Presented Posters

Mass Spectrometric Assessment of Carcinogen Exposure in Populations

E. Bailey, A. G. Brooks, P. B. Farmer, and B. Street. Monitoring exposure to 4,4′-methylene-bis(2-chloroaniline) through the gas chromatography–mass spectrometry measure of adducts to hemoglobin 175

M. D. Friesen, L. Garren, J.-C. Bereziat, F. Kadlubar, and D. Lin. Gas chromatography–mass spectrometry analysis of 2-amino-1-methyl-6-phenylimidazo [4,5-*b*] pyridine in urine and feces 179

K. W. Turteltaub, J. S. Vogel, C. Frantz, M. H. Buonarati, and J. S. Felton. Low-level biological dosimetry of heterocyclic amine carcinogens isolated from cooked food 183

G. Schäffler, C. Betz, and E. Richter. Mass spectrometric analysis of tobacco-specific hemoglobin adducts 187

J. O. Lay, Jr., M. P. Chiarelli, M. S. Bryant, and R. W. Nelson. Detection and characterization of DNA adducts at the femtomole level by desorption ionization mass spectrometry 191

Assessment of Carcinogen Exposure and the Use of Other Risk Markers in Populations

J. L. Autrup, J. Schmidt, and H. Autrup. Exposure to aflatoxin B_1 in animal-feed production plant workers 195

R. M. Santella, Y-J. Zhang, C-J. Chen, L-L. Hsieh, C-S. Lee, B. Haghighi, G-Y. Yang, L-W. Wang, and M. Feitelson. Immunohistochemical detection of aflatoxin B_1–DNA adducts and hepatitis B virus antigens in hepatocellular carcinoma and nontumorous liver tissue 199

S. G. Carmella, S. S. Kagan, and S. S. Hecht. Evidence that a hemoglobin adduct used for dosimetry of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone is a carboxylic ester 203

N. J. Gorelick and N. L. Reeder. Detection of multiple polycyclic aromatic hydrocarbon–DNA adducts by a high-performance liquid chromatography–^{32}P-postlabeling method 207

G. Sabbioni. Hemoglobin binding of aromatic amines: molecular dosimetry and quantitative structure–activity relationships for *N*-oxidation 213

N. J. van Sittert, G. D. J. Beulink, E. W. N. van Vliet, and H. van der Waal. Monitoring occupational exposure to ethylene oxide by the determination of hemoglobin adducts 217

C. J. Calleman, E. Bergmark, L. G. Stern, and L. G. Costa. A nonlinear dosimetric model for hemoglobin adduct formation by the neurotoxic agent acrylamide and its genotoxic metabolite glycidamide 221

J. Gallagher, M. George, M. Kohan, C. Thompson, T. Shank, and J. Lewtas. Detection and comparison of DNA adducts after *in vitro* and *in vivo* diesel emission exposures 225

C. Hansen, I. Asmussen, and H. Autrup. Detection of carcinogen-DNA adducts in human fetal tissues by the ^{32}P-postlabeling procedure 229

R. Mustonen, A. Försti, P. Hietanen, and K. Hemminki. Development of ^{32}P-postlabeling assay for 7-methyl guanines in human DNA 233

G. Assennato, G. M. Ferri, M. S. Tockman, M. C. Poirier, B. Schoket, A. Porro, V. Corrado, and P. T. Strickland. Biomarkers of carcinogen exposure and cancer risk in a coke plant 237

W. J. Bodell, G. Levay, and K. Pongracz. Investigation of benzene–DNA adducts and their detection in human bone marrow 241

E. Eder, C. Hoffman, S. Sporer, and S. Scheckenbach. Biomonitoring studies and susceptibility markers for acrolein congeners and allylic and benzyl compounds 245

S. De Flora, A. Izzotti, F. D'Agostini, G. A. Rossi, and R. M. Balansky. Pulmonary alveolar macrophages in molecular epidemiology and chemoprevention of cancer 249

C. G. Gairola, H. Wu, R. C. Gupta, and J. N. Diana. Mainstream and sidestream cigarette smoke-induced DNA adducts in C57Bl and DBA mice 253

A. Weston, E. D. Bowman, P. G. Shields, G. E. Trivers, M. C. Poirier, R. M. Santella, and D. K. Manchester. Detection of polycyclic aromatic hydrocarbon–DNA adducts in human lung 257

V. L. Wilson, B. G. Taffe, P. G. Shields, A. C. Povey, and C. C. Harris. Detection and quantification of 8-hydroxy-deoxyguanosine adducts in peripheral blood of people exposed to ionizing radiation 261

N. Rothman, M. C. Poirier, R. A. Haas, A. Correa-Villasenor, P. Ford, J. A. Hansen, T. O'Toole, and P. T. Strickland. Association of PAH–DNA adducts in peripheral white blood cells with dietary exposure to polyaromatic hydrocarbons 265

H-I. Kang, C. Konishi, T. Kuroki, and N-K. Huh. A highly sensitive and specific method for quantitation of *O*-alkylated DNA adducts and its application to the analysis of human tissue DNA 269

S. J. Culp, M. C. Poirier, and F. A. Beland. Biphasic removal of DNA adducts in a repetitive DNA sequence after dietary administration of 2-acetylaminofluorene 273

B. A. Smith, N. F. Fullerton, A. Aidoo, R. H. Heflich, and F. A. Beland. DNA adduct formation in relation to lymphocyte mutations and lung tumor induction in F344 rats treated with the environmental pollutant 1,6-dinitropyrene 277

G. Birner, S. Vamvakas, W. Dekant, and D. Henschler. Nephrotoxic and genotoxic *N*-acetyl-*S*-dichlorovinyl-*L*-cysteine is a urinary metabolite after occupational 1,1,2-trichloroethene exposure in humans: implications for the risk of trichloroethene exposure 281

M. Vanderlaan, M. Hwang, and T. Djanegara. Immunoaffinity purification of dietary heterocyclic amine carcinogens 285

G. Talaska, M. Schamer, P. Skipper, S. Tannenbaum, N. Caporaso, F. Kadlubar, H. Bartsch, and P. Vineis. Carcinogen-DNA adducts in exfoliated urothelial cells: techniques for noninvasive human monitoring 289

S. A. Bingham, A. B. Shah, A. Ellul, J. H. Cummings, and I. K. O'Neill. Magnetic microcapsules as novel biomonitors of cross-linking agents and diet-dependent reactive oxygen species in the human gastrointestinal tract 293

P. J. E. Compton-Quintana, R. H. Jensen, W. L. Bigbee, S. G. Grant, R. G. Langlois, M. T. Smith, and S. M. Rappaport. Use of the glycophorin A human mutation assay to study workers exposed to styrene 297

R. J. Šrám, B. Binková, L. Dobiáš, P. Rössner, J. Topinka, D. Veselá, D. Veselý, J. Stejskalová, H. Bavorová, and V. Řeřicha. Monitoring genotoxic exposure in uranium miners 303

B. Schoket, D. H. Phillips, M. C. Poirier, and I. Vincze. DNA adducts in peripheral blood lymphocytes from aluminum production plant workers determined by ^{32}P-postlabeling and enzyme-linked immunosorbent assay 307

Contributed Articles

G. C. Hard, I. S. Rogers, K. P. Baetcke, W. L. Richards, R. E. McGaughy, and L. R. Valcovic. Hazard evaluation of chemicals that cause accumulation of $\alpha_{2\mu}$-globin, hyaline droplet nephropathy, and tubule neoplasia in the kidneys of male rats. 313

D. J. O'Brien, J. B. Kaneene, and R. Poppenga. The use of mammals as sentinels for human exposure to toxic contaminants in the environment 351

R. G. Wones, C. C. Deck, B. Stadler, S. Roark, E. Hogg, and L. A. Frohman. Effects of drinking water monochloramine on lipid and thyroid metabolism in healthy men. 369

R. G. Wones, C. C. Deck, B. Stadler, S. Roark, E. Hogg, and L. A. Frohman. Lack of effect of drinking water chlorine on lipid and thyroid metabolism in healthy humans. 375

E. S. Johnson. Important aspects of the evidence for TCDD carcinogenicity in man. 383

Announcement 391

Instructions to Authors 393

Published Volumes of EHP 395